



BEETUP FLO®

HERBICIDE

PCS 05272

**FOR USE AS AN AGRICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY**

A suspension concentrate (SC) containing 160 g/L (16% w/w) phenmedipham. Also Contains 1,2-benzisothiazolin-3-one.
For control of broad-leaved weeds in sugar beet, fodder beet, red beet and mangels.

SAFETY INFORMATION WARNING

May cause an allergic skin reaction.

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Avoid breathing spray.


Wear protective gloves/protective clothing/eye protection/face protection.

If skin irritation or rash occurs: Get medical advice/attention.

If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 05272

SHAKE WELL BEFORE USE
PROTECT FROM FROST

For Batch Number and Manufacturing Date see container.

Authorisation Holder: UPL Holdings Coöperatief U.A.,
Claudius Prinsenlaan 144 A, Blok A, 4818CP Breda,
The Netherlands

Marketing Company: UPL Europe Ltd

Engine Rooms (1st Floor), Birchwood Park, Warrington, WA3 6YN, UK
Telephone: +44 (0)1925 819999 Fax: +44 (0)1925 817425

For 24 hour emergency information contact:
CARECHEM24 : +44 (0)1235 239670

5L



Crops/situations	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha/crop)	Latest time of application
Sugar beet, fodder beet, red beet and mangles	3.0	6.0	Before crops meet between the rows

SAFETY PRECAUTIONS

OPERATOR PRECAUTIONS

OPERATORS MUST WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE WATER WITH THE PRODUCT OR ITS CONTAINER (DO NOT CLEAN APPLICATION EQUIPMENT NEAR SURFACE WATER/ AVOID CONTAMINATION VIA DRAINS FROM FARMYARDS AND ROADS).

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

STORAGE AND DISPOSAL

KEEP AWAY FROM FOOD, DRINK OR ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rising device.

After use, thoroughly wash out the sprayer with a minimum of two flushes of water. Add washings to sprayer at time of filling and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS/WARNINGS

1. The application of BEETUP FLO to crops suffering from stress may lead to a check to growth from which the crops may not recover and in some cases crop losses may occur.
2. If the temperature is 21°C or is likely to rise to this temperature for the period of twenty-four hours after application do not spray as serious crop damage can occur.
3. If substantial day to night fluctuations in temperature occur shortly before or after the application crop growth may be checked.
4. Weeds are best controlled at the seeding stage when growing actively in warm, moist, sunny conditions. Under poor growing conditions control may be reduced.
5. BEETUP FLO will not control perennial weeds.
6. DO NOT apply to wet foliage or during rainfall or drizzle.
7. DO NOT overlap spray bouts.
8. DO NOT apply any of the residual herbicide mixtures more than three times.
9. DO NOT apply a residual herbicide mixture to sand soils.
10. DO NOT roll or harrow the crop within seven days before or after applying BEETUP FLO.
11. Avoid spray drift onto neighbouring crops.

CROP SPECIFIC INFORMATION

WEEDS CONTROLLED

BEETUP FLO should be used as a repeat low dose treatment when weeds are still small. Best control is achieved at the cotyledon stage. It is normal for a repeat low dose treatment to be applied as an overall spray.

BEETUP FLO is a contact herbicide that is dependent for action on light and temperature. This affects both weed control and crop safety. It is therefore important that all instructions in this label are read and followed. This is particularly important for application under hot conditions, to crops under stress and for tank mixtures.

BEETUP FLO is absorbed through the leaves of the weeds and effects should be seen within five to six days of spraying. Weed response will vary according to conditions at spraying and the stage of growth of individual weeds. Weeds are best controlled at the seeding stage when growing actively in warm, moist, sunny conditions. Under poor growing conditions control may be reduced.

Susceptible weeds quickly die and in warm weather will be dead in 3-4 days. More tolerant weeds may take longer to die. Under cool conditions results may not be complete for as long as 7-10 days. BEETUP FLO will not prevent germination of weed seed after treatment and where this occurs further applications may be made provided the crop is in a healthy state and other conditions for safe use are complied with.

As BEETUP FLO has only contact action it is not affected by soil type and can be used on organic and peaty soils as well as mineral soils.

Susceptibility of weeds at cotyledon stage 3 x 2 l/ha

Annual meadow-grass	R
Black bindweed	MS
Common chickweed	S
Fat-hen	MS
Field pansy	MR
Field penny cress	-
Knotgrass	MS
Scentless mayweed	MR
Redshank	-

Susceptibility of weeds at cotyledon stage 2 x 3 l/ha

R
MS
MS
MS
MS
MS
R
-
R

S = susceptible, MS = moderately susceptible, MR = moderately resistant, R = resistant.

TREATMENT OF CROPS

CROP TOLERANCE

All varieties of sugar beet, fodder beet and mangels can be sprayed at any stage of growth if healthy, however, **do not** apply to red beet before the crop has reached the fully expanded cotyledon stage. Timing should be determined by the stage of growth of the weeds present. The safety of BEETUP FLO to the beet crop will vary with conditions around the time of application. As the beet plants are small and grow rapidly they can be affected by almost any change in conditions and can become stressed. This may not always be visible until affected by other factors such as herbicide spray.

The application of BEETUP FLO to crops suffering from stress may lead to a check to growth from which the crops may not recover and in some cases crop losses may occur. The most important causes of stress are listed below:

- **High Temperatures:**

If the temperature is 21°C or is likely to rise to this temperature for the period of twenty-four hours after application, do not spray as serious crop damage can occur. Where temperatures have been as high as 21°C and are falling, BEETUP FLO may be applied at a reduced rate but some reduction in weed control should be expected. This is particularly useful in the evening after a hot day. A return to temperatures of 21°C and

above in the following day may cause some check to the crop but this will normally be outgrown. If substantial day to night fluctuations in temperature occur shortly before or after the application crop growth may be checked.

- **High Light Intensity:**

As BEETUP FLO is activated by light, bright cloudless days may cause BEETUP FLO to check the growth of the crop especially if it is already under another stress.

- **Nutrient Deficiency:**

Crop suffering from nutrient deficiencies such as nitrogen, copper, manganese, magnesium or lime may be adversely affected when sprayed with BEETUP FLO. Growers should ensure that nutrient status of the soil is correct before the crop is drilled.

- **Effects of Other Herbicides:**

Damage or stress from pre-planting or pre-emergence herbicides can be increased by application of BEETUP FLO. When such damage or stress is apparent allow the symptoms to disappear before applying the BEETUP FLO.

- **Damage from Insect, Other Pests or Fungal Attack:**

This will reduce the plants tolerance to BEETUP FLO and the effects of such pest attack as mangel fly should be alleviated before application of BEETUP FLO.

- **Frost:**

Plants treated with BEETUP FLO may be more affected by frost following treatment than untreated crops.

- **Hail or Wind:**

Plants suffering from hail damage or wind damage due to either rocking in the soil or from wind borne soil particles should not be treated until the effects of this damage have been outgrown.

REPEAT LOW DOSE PROGRAMME

This programme depends on the application at a very early stage of weed growth. This can then be followed with repeat doses at the same rate of use as further weeds appear. Timing is very important and great care must be exercised in monitoring the appearance of weed seedlings so that they do not become too large.

- **Timing**

Apply when the weeds have their cotyledons fully expanded. In a mixed weed population, timing should be determined by the most forward plants. Where occasional isolated weeds become established ahead of normal weed populations they will generally be held in check by a programme of treatment. Spray whenever any weeds reach the stage of growth above.

- **Rates of Use**

2 litres BEETUP FLO per hectare for a treatment programme of 3 applications or 3 litres per hectare for a treatment of 2 applications. Ideally apply BEETUP FLO in tank mix with other sugar beet herbicide products as part of a low volume programme. Supported tank mixes can be found on the UPL website www.upleurope.com.

- **Application**

Apply overall as a FINE spray (BCPC definition). Application must only be 80-100 litres of water per hectare. Even cover with a fine spray is essential for good results. The water volumes and nozzles used should be selected to achieve this aim. Do not use water volume in excess of 100l/ha as nozzle blockage may occur.

MIXING

Half fill the spray tank with water and commence agitation. Add the tank-mix partner first and allow complete dispersion. Add the required amount of BEETUP FLO to the sprayer tank and fill up with water whilst agitating. Use immediately. Continue agitation until spraying is complete.

Application should be completed within two hours of mixing as there is a risk of nozzle blockage, especially if the water temperature is lower than 5°C.

Do not tank mix BEETUP FLO with manganese sulphate.

COMPATABILITY

The effectiveness of BEETUP FLO can be extended by tank-mixing with other herbicides. These mixtures must be sprayed overall and form part of a low volume programme of treatments. Only recommended mixtures should be made and manufacturer's recommendations consulted before mixing.

The UPL website (www.upl-ltd.com/ie) provides further information on tank mixtures.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable. However, as we cannot exercise control over their storage, handling, mixing, use or of the weather conditions before, during and after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

Brand names used in this label may be registered trademarks of UPL Europe Ltd, or other manufacturers in which propriety rights may exist.

To access the Safety Data Sheet for BEETUP FLO[®] please use the link below or scan the code.



www.upl-ltd.com/ie/product-details/beetup-flo

Alternatively, contact your local supplier.



BEETUP FLO®

HERBICIDE

PCS 05272

**FOR USE AS AN AGRICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY**

A suspension concentrate (SC) containing 160 g/L (16% w/w) phenmedipham. Also Contains 1,2-benzisothiazolin-3-one.
For control of broad-leaved weeds in sugar beet, fodder beet, red beet and mangels.

SAFETY INFORMATION WARNING

May cause an allergic skin reaction.

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Avoid breathing spray.


Wear protective gloves/protective clothing/eye protection/face protection.

If skin irritation or rash occurs: Get medical advice/attention.

If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 05272

SHAKE WELL BEFORE USE
PROTECT FROM FROST

For Batch Number and Manufacturing Date see container.

Authorisation Holder: UPL Holdings Coöperatief U.A.,
Claudius Prinsenlaan 144 A, Blok A, 4818CP Breda,
The Netherlands

Marketing Company: UPL Europe Ltd

Engine Rooms (1st Floor), Birchwood Park, Warrington, WA3 6YN, UK
Telephone: +44 (0)1925 819999 Fax: +44 (0)1925 817425

For 24 hour emergency information contact:
CARECHEM24 : +44 (0)1235 239670

5L

October 2020

