



ADAMA FALCON[®] RATE CARD

Grass weed	Weed growth stage	Rate (L/ha)
Volunteer wheat	Optimum: 2 leaves unfolded to end of tillering. Latest: Stem erect	0.5 – 0.7
Volunteer barley	Optimum: 2 leaves unfolded to end of tillering. Latest: Stem erect	0.3 – 0.7
Sterile brome	Optimum: 2 expanded leaves to fully tillered	0.5 – 0.75
Barley cover crops	Optimum: 2 leaves unfolded to stem erect Latest: 2nd node detectable	0.7 – 1.0
Wild oat	Optimum: 2 leaves unfolded to early tillering Latest: Stem erect	0.5 – 0.7
Common couch	3 leaves unfolded (when majority of shoots have emerged and are approx. 15 cm tall)	1.0 – 1.5
Rye grass (from seed)	Optimum: 2 leaves unfolded to early tillering Latest: Before stem erect	1.2
Black-grass	Optimum: 2 leaves unfolded to stem erect Latest: Booting	0.7 – 1.0

- To avoid the build-up of resistance do not apply products containing ACCase inhibitor herbicide more than twice to any crop.
- Speed of kill will be more rapid when weeds are actively growing under warm conditions with adequate moisture.

Use plant protection products safely. Always read the label and product information before use. For further product information, including warning phrases and symbols, refer to www.adama.com/uk or call The Technical Helpline on 01635 876 622.

ADAMA Agricultural Solutions UK Ltd. Third Floor East, 1410 Arlington Business Park, Theale, Reading RG7 4SA. Telephone 01635 860 555. UKenquiries@adama.com.