



Manganese 64

Croplink Manganese 64 contains:			
	% w/v	% w/w	g/l
Manganese (Mn)	7.6	6.4	64

Complexed with carboxylic acid and EDTA

EEC FERTILISER. Use only where there is a recognised need. Do exceed the appropriate application rates

CEREALS

At the first sign of deficiency apply Croplink Liquid Manganese at 3 l/ha regardless of the crop stage. Repeat applications may be necessary. When applications are planned into a crop protection programme, or where crops are at high risk from sub-clinical deficiencies, the following timings will ensure adequate levels:

WINTER CEREALS

1.5 l/ha at Growth Stage 4-6 (Zadoks 30-31). Pseudo stem erect to 1st node detectable. plus

1.5 l/ha at Growth Stage 8 (Zadoks 37). Flag leaf just visible.

For early drilled winter cereals on susceptible soil types an autumn application at 1.5 l/ha may be applied.

SPRING CEREALS

1.5 l/ha at the 3-5 leaf stage plus

1.5 l/ha at the Growth Stage 7-8 (Zadoks 32-37) 2nd node to flag leaf just visible.

SUGAR BEET

At the first sign of deficiency apply 3 l/ha regardless of crop stage. Repeat if necessary.

On susceptible soil types maximum benefit will be obtained from planned applications made before deficiency develops.

EARLY APPLICATION

1.5 l/ha When there is enough leaf to absorb the spray. This often coincides with early post-emergence weed control.

plus 1.5 l/ha When the crop has 4-6 leaves.

MID-SEASON

3 l/ha In high risk situations when full leaf cover is obtained.

SAFETY PRECAUTIONS.

R48 RISK OF SERIOUS DAMAGE TO HEALTH BY PROLONGED EXPOSURE BY INHALATION AND IF SWALLOWED.

R51/53 TOXIC TO AQUATIC ORGANISMS. MAY CAUSE LONG TERM DAMAGE IN THE AQUATIC ENVIRONMENT

S(2-) KEEP OUT OF REACH OF CHILDREN.

S23 DO NOT BREATHE SPRAY



Net Contents: 10 litres

Croplink (Ireland) Ltd. , Springhill, Carrigtwohill, Co. Cork T45 F685

Tel: 021 488 2806