



NETTLES



DOCKS

CROP
LINK

SCRUBKILLER

ULTRA



Ultimate spot weed
control in Grassland

Fully approved for handheld spot treatment

Targets the roots for long term control



GORSE



THISTLES



BRAMBLES

PRODUCT INFORMATION

- **PCS Number:** 06734
- **Active Ingredients:** 240g/L Triclopyr + 60g/L Clopyralid
- **Pack Size:** 1L + 3L
- **Formulation:** Emulsifiable Concentrate
- **Recommended Dose:** 60ml Scrubkiller Ultra in 10L water
- **Maximum Total Dose:** Once per year
- **Latest Timing:** Seven days before grazing. Apply between 1st March and 31st October

BEST PRACTISE GUIDELINES

- For use only in established grassland >12 months
- Users should be reminded to wear appropriate PPE: face-shield, gloves, coveralls and rubber boots
- Spray to thoroughly wet the foliage of the plant but run-off should be avoided
- If weeds grow beyond ideal stage, consider cutting the sward and spraying the re-growth
- Allow 2-3 weeks after cutting before applications of Scrubkiller Ultra are made to allow sufficient re-growth
- Grazing can resume 7 days after application. This may need to be longer if foliage of poisonous weeds is present
- Allow an interval of minimum 21 days, ideally 28 days before cutting
- Scrubkiller Ultra should not be tank-mixed with phenoxy herbicides e.g. MCPA, 2,4-D, CMPP – this may compromise translocation of Scrubkiller Ultra to the roots
- Keep pets out of treated areas until the spray has dried (normally about 2 hours)
- If weed populations exceed 5% then a broad-acre treatment with Pasture Pack may be more cost effective

KEY TIMINGS FOR BEST CONTROL

Weed	Timing
Docks	Before plant reaches 20 cm, or after start of regrowth post cutting or topping
Nettles	About 25 cm tall, prior to flowering
Thistles	Before flowering with adequate foliage present when plants are no more than 8-10 cm
Gorse	June – August pre-senescence
Brambles	June – August pre-senescence

WEED CONTROL RATINGS

Note weed control of non-label weeds below based on limited data and therefore the user bears the risk as actual results may vary.

Annual Weeds	Control
Bindweed (black)	★★★
Bindweed (field)	★★★
Bristly ox-tongue	★★
Chickweed	★★★
Corn chamomile	★★★
Corn marigold	★★★
Cranesbill	★★★
Dead-nettles	★★★
Fat-Hen (2TL)	★★★
Fool's parsley	★★★
Groundsel	★★★
Himalayan balsam	★★★
Knotgrass (4TL)	★★★
Medick	★★★
Mayweeds	★★★
Nightshade (black)	★★★
Pale persicaria	★★★
Scarlet pimpernel	★★
Shepherd's purse	★★★
Speedwells	★
Yellow rattle	★★

Perennial Weeds	Control
Bramble	★★★
Broom	★★★
Buttercups	★
Clover, trefoil	★★★
Coltsfoot	★★★
Cow parsley	★★
Daisy, common	★★
Dandelion	★★
Docks	★★★
Gorse	★★★
Ground Elder	★★★
Hawthorn	★★★
Hemlock	★★★
Hogweed (giant)	★★★
Horsetail (Equisetum)	★
Japanese Knotweed	★★★
Knapweed (common)	★★
Lesser celandine	★★★
Mallow	★★★
Mugwort	★★★
Nettle (common)	★★★
Plantain (greater)	★★
Plantain (ribwort)	★★★
Ragwort	★
Rosebay Willowherb	★★★
Self-heal	★★★
Silverweed	★★★
Sorrel (common)	★★★
Thistles	★★★
Vetch, tare	★★★
Yarrow	★★
Yellow/Flag Iris	★★★



Key

- ★★★ Good Control
- ★★ Moderate Control
- ★ Some Control

The Stewardship logo, featuring a stylized flame or leaf icon to the left of the word 'Stewardship' in a large, bold, white font. Below it, the tagline 'Protect a better tomorrow' is written in a smaller, white font.

Pesticides are soluble in water, extra care must be taken to prevent pollution. It is important to adopt effective Stewardship and observe best practice guidelines.

- Keep pesticides in a locked, bunded store.
- Application must be made by a qualified operator.
- Check and calibrate equipment regularly.
- Clear up any spills immediately
- Consider weather and soil conditions before application.
- Never exceed label rates.
- Clean and wash sprayers at the end of the day.
- Dispose of washings on an untreated area of grassland.



Croplink (Ireland) Ltd,
Springhill
Carrigtwohill
Co. Cork,
T45 F685



info@croplink.ie



www.croplink.ie



Office Tel.	021 4882806
Office contact:	Theresa McGeeney
Thomas Delaney	087 2553564
Eddie Young	087 1056944
Scott Lovell	087 6379079