



Zinc 93

Crop Link Zinc 93 contains 9.3% W/V Zinc in solution chelated with E.D.T.A.

FOLIAR APPLICATION Of Croplink Zinc : Rates & Timings

At first sign of zinc deficiency apply Croplink Zinc at 3 ltr/ha regardless of crop stage. Repeat applications may be necessary.

When applications are planned into a crop protection programme or where crops are at high risk from sub-clinical deficiencies, the following rates and timings will ensure adequate levels.

WINTER CEREALS

Timing

A planned application programme is critical. Autumn application is important on sites with a history of zinc deficiency.

Apply Croplink Zinc 93 at	1.5 litres/ha	Autumn/Winter
plus Croplink Zinc 93 at	1.5 litres/ha	Growth stage 4-6 (Zadoks 30-31)
plus Croplink Zinc 93 at	1.5 litres/ha	Growth stage 8 (Zadoks 37)

SPRING CEREALS

Apply Croplink Zinc 93 at	1.5 litres/ha	at 3-5 leaf stage
plus Croplink Zinc 93 at	1.5 litres/ha	Growth stage 7-8 (Zadoks 32-37)



SAFETY PRECAUTIONS.

R36/38 IRRITATING TO EYES AND SKIN.
S(2-) KEEP OUT OF REACH OF CHILDREN.
S25 AVOID CONTACT WITH THE EYES.
PROTECT FROM FROST.

Net Contents: 10 litres

**Croplink (Ireland) Ltd. , Springhill, Carrigtwohill, Co. Cork T45 F685
Tel: 021 488 2806**